

Intersexuality and the Cricket Frog Decline: Historic and Geographic Trends

Amy L. Reeder,¹ Marilyn O. Ruiz,² Allan Pessier,³ Lauren E. Brown,⁴ Jeffrey M. Levensgood,⁵ Christopher A. Phillips,⁷ Matthew B. Wheeler,¹ Richard E. Warner,⁶ and Val R. Beasley⁵

¹Department of Animal Sciences, and ²Department of Veterinary Pathobiology, University of Illinois at Urbana-Champaign, Champaign, Illinois, USA; ³University of Illinois Zoological Pathology Program, Loyola University Medical Center, Maywood, Illinois, USA; ⁴Department of Biological Sciences, Illinois State University, Normal, Illinois, USA; ⁵Department of Veterinary Biosciences, and ⁶Department of Natural Resources and Environmental Sciences, University of Illinois at Urbana-Champaign, Champaign, Illinois, USA; ⁷Center for Biodiversity, Illinois Natural History Survey, Champaign, Illinois, USA

Exposure to anthropogenic endocrine disruptors has been listed as one of several potential causes of amphibian declines in recent years. We examined gonads of 814 cricket frogs (*Acris crepitans*) collected in Illinois and deposited in museum collections to elucidate relationships between the decline of this species in Illinois and the spatial and temporal distribution of individuals with intersex gonads. Compared with the preorganochlorine era studied (1852–1929), the percentage of intersex cricket frogs increased during the period of industrial growth and initial uses of polychlorinated biphenyls (PCBs) (1930–1945), was highest during the greatest manufacture and use of *p,p*-dichlorodiphenyltrichloroethane (DDT) and PCBs (1946–1959), began declining with the increase in public concern and environmental regulations that reduced and then prevented sales of DDT in the United States (1960–1979), and continued to decline through the period of gradual reductions in environmental residues of organochlorine pesticides and PCBs in the midwestern United States (1980–2001). The proportion of intersex individuals among those frogs was highest in the heavily industrialized and urbanized northeastern portion of Illinois, intermediate in the intensively farmed central and northwestern areas, and lowest in the less intensively managed and ecologically more diverse southern part of the state. Records of deposits of cricket frog specimens into museum collections indicate a marked reduction in numbers from northeastern Illinois in recent decades. These findings are consistent with the hypothesis that endocrine disruption contributed to the decline of cricket frogs in Illinois. **Key words:** *Acris crepitans*, amphibian, cricket frogs, endocrine disruption, environmental contaminants, Illinois, intersexuality. *Environ Health Perspect* 113:261–265 (2005). doi:10.1289/ehp.7276 available via <http://dx.doi.org/> [Online 7 December 2004]

Amphibian declines have been documented in many parts of the world (Alford et al. 2001; Blaustein and Wake 1990; Houlihan et al. 2000; Tyler 1991), which is of concern because these species are important grazers, prey species, and predators in aquatic and terrestrial ecosystems and serve as valuable sentinels of ecologic integrity. For many amphibian species, including the cricket frog *Acris crepitans*, the causes of declines are unclear. This species is indigenous to the eastern half of the United States and has experienced a marked decline in portions of its range in the last 25 years (Brodman and Kilmurry 1998; Caspar 1998; Jung 1993; Mierzwa 1998; Minton 1998; Moriarity 1998; Mossman et al. 1998). In 1961, it was the most common amphibian in Illinois (Smith 1961), and it is still abundant in southern Illinois. However, cricket frogs were rarely encountered in amphibian surveys conducted in portions of northern Illinois in recent decades (Ludwig et al. 1992; Vogt 1981; Phillips CA, Brown LE, personal observation).

A variety of industrial compounds and by-products disrupt endocrine function, and exposure to such chemicals may contribute to amphibian declines. Polychlorinated biphenyls (PCBs), polychlorinated dibenzofurans (PCDFs), and polychlorinated dibenzo-*p*-dioxins (PCDDs) may produce estrogenic, antiestrogenic, and antiandrogenic effects

(Jansen et al. 1993; Krishnan and Safe 1993; Li and Hansen 1996; Li et al. 1994; Malby et al. 1992). For example, PCBs may affect sexual differentiation in the slider turtle *Trachemys scripta* (Bergeron et al. 1994), various frogs (Qin et al. 2003), and fish (Matta et al. 1998). Moreover, our laboratory associated exposures of cricket frog tadpoles to antiestrogenic PCBs and PCDFs with marked masculinization of sex ratios (Reeder et al. 1998). In addition, polycyclic aromatic hydrocarbons (PAHs) from coal tar and smoke from combustion of coal, oil, gas, wood, and garbage are widely disseminated endocrine disruptors (Chaloupka et al. 1992, 1993; Machala et al. 2001; ATSDR 1995; Thomas and Budiantara 1995).

Pesticides now banned from the market and compounds still sold can adversely affect animal reproduction. For example, the organochlorine pesticide *p,p*-dichlorodiphenyltrichloroethane (DDT) and its metabolites/environmental products have demasculinized birds (e.g., gulls, *Larus californicus* and *L. occidentalis*), reptiles (*Alligator mississippiensis*), and fish (Baatrup and Junge 2001; Fry and Toone 1981; Guillette et al. 1994, 1995; Metcalfe et al. 2000). Also, DDT induced vitellogenin production in male African clawed frogs (*Xenopus laevis*) and slider turtles (*T. scripta*), and the organochlorine insecticides toxaphene and dieldrin induced this protein in male

X. laevis (Palmer and Palmer 1995; Palmer et al. 1998). Furthermore, there is concern that the widely used herbicide atrazine may impair reproduction and/or development of amphibians. At high concentrations, atrazine decreased growth in gray treefrogs (*Hyla versicolor*) and increased time to metamorphosis in *X. laevis* (Diana et al. 2000; Sullivan and Spence 2003). At much lower concentrations, atrazine in combination with nitrate reduced growth of *X. laevis* (Sullivan and Spence 2003). Hayes et al. (2002, 2003) associated hermaphroditism and gonadal dysgenesis in amphibians with very low aquatic concentrations of atrazine (0.1 µg/L), whereas Carr et al. (2003) observed significantly increased intersexuality only at higher concentrations (25 µg/L).

Although it is apparent that exposure to anthropogenic compounds may harm amphibians through changes in functional sex ratios, reduced size at metamorphosis, and delayed maturation, the distribution of intersex frogs geographically and historically remains to be

Address correspondence to V.R. Beasley, Department of Veterinary Biosciences, College of Veterinary Medicine, University of Illinois, 2001 S. Lincoln Ave., Urbana, IL 61802 USA. Telephone: (217) 333-9360. Fax: (217) 244-1652. E-mail: val@uiuc.edu

We thank M. Post for technical assistance. We greatly appreciate the following individuals and their respective institutions for allowing us to examine cricket frogs in their natural history collections: H. Voris and A. Resetar, Field Museum of Natural History; R. Axtell, Department of Biological Sciences, Southern Illinois University; E. Moll, Department of Zoology, Eastern Illinois University; R. Brandon, Department of Zoology, Southern Illinois University; K. Cummings, Illinois Natural History Survey; the Department of Biological Sciences, Illinois State University; R.P. Reynolds and R. McDiarmid, National Museum of Natural History, Smithsonian Institution; G. Thurow, Department of Biology, Western Illinois University; J.R. Purdue and D. Bakken, Illinois State Museum; G. Schneider, Museum of Zoology, University of Michigan; R. Vasile, M. Hennen, and S. Sullivan, Chicago Academy of Sciences; E. Censky, Carnegie Museum of Natural History; J. Cadle, Academy of Natural Sciences; and J.W. Wright, Natural History Museum of Los Angeles County.

Funding was provided by the John G. Shedd Aquarium through support from the Dr. Scholl Foundation.

The authors declare they have no competing financial interests.

Received 24 May 2004; accepted 7 December 2004.

characterized. In this study, we examined gonads of *A. crepitans* from museum specimens to elucidate relationships between the decline of this species and the temporal and spatial occurrence of intersexuality.

Materials and Methods

Natural history museums are valuable resources for estimations of species distributions and health status over time (Shaffer et al. 1998). We examined specimen records from 16 museums to determine where and when cricket frogs were collected in Illinois. To determine whether cricket frogs were not available because collecting was not conducted, we compared cricket frog records with those of all anuran collections from the state. Our rationale was that scientists collecting anurans and placing them in museums as voucher specimens would not consistently preclude cricket frogs.

Museum specimens from throughout Illinois (Figure 1) were examined to compare cricket frog gonadal sex in three regions of the state during five time periods. The three regions are distinguished by human population density and physiographic characteristics. The northeast region includes 11 counties with high human population densities in and surrounding the Chicago metropolitan area. The central band of 66 counties, which was formerly largely prairie, is dominated by low topographic relief, fertile soils, intensive maize and soy agriculture, and low human population density. The southernmost region

includes 25 counties with mixed crops, pastures, and wooded hills as well as low human population density. The five time periods studied included *a*) a preorganochlorine period (1852–1929); *b*) an era of PCB use and industrialization that predates use of DDT (1930–1945); *c*) a period of rapidly increasing DDT use and further industrialization (1946–1959); *d*) a period of declining use and then a ban on sales of DDT as well as initial

measures to limit pollution from industries (1960–1979); and *e*) a period associated with a substantial decline in environmental residues of organochlorine insecticides and other persistent halogenated organic air and water pollutants in the Midwest (1980–2001).

The gonads of 814 cricket frogs were examined *in situ* with a dissecting microscope to identify sex. Because there is no evidence that testicular tissue develops within the




Figure 2. (A) Numbers of cricket frog specimens from Illinois deposited in museum collections relative to numbers of other anurans in museums collected in the state from 1852 to 2001. (B) Numbers of cricket frog specimens from northeastern Illinois deposited in museum collections relative to numbers of other anurans in museums collected in that region from 1852 to 2001.




Figure 1. Number by county of cricket frogs (*n* = 814) collected in Illinois from 1852 to 1996 and examined for gonadal sex determination.




Figure 3. Distributions of total numbers of Illinois cricket frogs in museum collections for the five time periods. Each circle represents one museum specimen from that county.

female tract, females were identified by the presence of oocytes and excluded from further analysis. Testes, intersex gonads, and poorly differentiated gonads were removed for histologic examination. When museums limited the proportion of cricket frogs from which gonads could be taken, we used a random numbers table to select a subset for histologic study. Collection dates, collecting localities, and notes from museum catalogs were recorded. Gonads were immersed in 70% ethanol, embedded in paraffin, and sectioned at 5 μ m. Depending on gonad size, longitudinal sections were obtained beginning 60–150 μ m from the outer edge. Two or three sections 60–120 μ m apart were selected for processing. Slides were stained with hematoxylin and eosin and examined with a light microscope for the presence of sperm production and oocytes. When oocytes were present within testicular tissue, they were relatively few in number, but each was several-fold larger than the spermatid ducts and thus took up a substantial percentage of testicular volume, making them easy to identify.

We used Systat 10.2 (SPSS, Inc., Chicago, IL) to construct contingency tables and perform Pearson chi-square analyses comparing proportions of male, female, and intersex individuals among the five time periods and three areas of Illinois. Further analysis incorporating temporal and spatial dimensions simultaneously was not possible because of the limited numbers of intersex specimens at some time points. Specifically, tables based on all five time periods and three regions broken down

by sex would have included cells with no data. The null hypotheses tested whether the proportion of male, intersex, and female frogs is the same for frogs captured during the five time periods and in the three areas of the state. We used the 5% significance level as the indicator of a statistically significant association.

Results

Records for all anurans totaled 12,661 specimens. Of these, 2,570 (20%) were *A. crepitans*, the frog species most often collected in Illinois. Years of collection ranged from 1852 to 2001. A trend of increasing numbers of frogs collected started in the late 1930s; there was a marked reduction during World War II, and then the rate of collecting markedly increased through the mid-1950s. Numbers of individual anurans collected declined sharply in the late 1950s, increased during the mid-1960s, and declined during the 1970s and 1980s. The numbers of frogs collected in Illinois have since increased. Cricket frog numbers were largely proportional to those of other anurans (Figure 2A). Other frog and toad species have been collected from the Chicago region in proportion to other regions of the state since 1960; however, few cricket frogs were obtained from that area in the same time frame (Figure 2B). Moreover, from 1980 to 2001, cricket frog collections remained high in the central and southern regions but declined even further around Chicago (Figure 3).

Intersexuality (hermaphroditism) was manifested in two forms: most intersex frogs

($n = 37$) had an ovotestis where proportionately large ova were present within testicular tissue, and a few ($n = 6$) had a complete testis and complete ovary. The proportions of specimens in each gonadal sex class differed significantly among the geographical regions in Illinois ($\chi^2 = 20.2$, 4 df, $p < 0.001$; Table 1, Figure 4A). Notably, in the urbanized northeastern portion of the state, the proportion of frogs that were intersex was much greater than in other areas, and the proportion of females was smaller than elsewhere. In southern Illinois, where agriculture and urbanization are least intensive, the proportion of intersex individuals was considerably smaller than in the other regions.

The proportion of specimens in each gonadal sex class differed markedly among the time periods of collection ($\chi^2 = 31.1$, 8 df, $p < 0.001$; Table 1, Figure 4B). From 1930 to 1945, the percentage of intersex individuals was notably increased, and from 1946 to 1959 it was greater than during any other time frame examined. Also, during 1946–1959, the proportion of females was reduced. During the most recent period, the proportion of intersexes was lowest of any period except for 1852–1929. In the 1990s, however, few cricket frogs were available from areas that previously had the most elevated intersex rates.

Discussion

Environmental contamination probably accounted for the historical and geographical trends in gonadal sex in Illinois cricket frogs and likely contributed to the decline of the species. In this research, it would not have been productive to assay contaminants in tissues of the frogs because of their small size, and because frogs are individually tagged but stored together in jars of fixative with conspecific individuals, which would enable post-mortem cross-contamination. Moreover, such use would consume the specimens, preventing any future examination for other research aims. Also, it would not be meaningful to assay contaminant mixtures at sites today because they would no longer be representative of what was present when the frogs were collected.

The absence of cricket frogs from northeastern Illinois in museum collections is consistent with reports indicating virtual disappearance of cricket frogs from this area (Ludwig et al. 1992; Phillips CA, Brown LE, personal observation; Vogt 1981). Heavy industrialization from the 1930s through the 1950s was accompanied by major releases of combustion products and organochlorine contaminants. Smokes contain abundant mixtures of PAHs that adsorb to particulates in air, soil, water, and sediment (Mumtaz and George 1995). Like coplanar PCDDs and similar organochlorines, some PAHs can act

Table 1. Numbers (percentages) of cricket frog specimens by gonadal sex and region and by gonadal sex and time period.

	Female	Intersex	Male	Total observed
Region				
Northeast	57 (32.8)	19 (10.9)	98 (56.3)	174
Central	146 (44.7)	16 (4.9)	165 (50.5)	327
South	138 (44.1)	8 (2.6)	167 (53.4)	313
Total	341 (41.9)	43 (5.3)	430 (52.8)	814
Time period				
1852–1929	44 (52.4)	1 (1.2)	39 (46.4)	84
1930–1945	35 (43.8)	6 (7.5)	39 (48.8)	80
1946–1959	46 (30.1)	17 (11.1)	90 (58.8)	153
1960–1979	76 (48.1)	10 (6.3)	72 (45.6)	158
1980–1996	140 (41.3)	9 (2.7)	190 (56.1)	339
Total	341 (41.9)	43 (5.3)	430 (52.8)	814




Figure 4. Deviations of observed from expected values of cricket frog sex (A) by time period and (B) by region. Expected values were determined from the overall data set using the chi-square test.

as antiestrogens (Chaloupka et al. 1992). By the 1930s, PCBs were being commercially produced (Hansen 1987), and coplanar PCBs and structurally similar PCDDs and PCDFs are potent antiestrogens. Thus, intensive industrial smoke emissions and commercial PCB production and widespread use coincided with the increase in the proportion of intersex cricket frogs. A possible outcome of exposure to various PAH and organochlorine antiestrogens is masculinization of juvenile cricket frogs and skewed sex ratios, as noted in the Chicago region in this study, which is similar to what we found previously at a hazardous waste site at Crab Orchard National Wildlife Refuge, where cricket frogs were contaminated with coplanar PCBs and PCDFs (Reeder et al. 1998).

The greatest proportions of intersex in cricket frogs of Illinois during 1946–1959 corresponded with a rapid increase in use of DDT in the United States (Mellanby 1992; U.S. Army Service Forces 1946). Large-scale DDT applications in Illinois for mosquito control began in 1945, followed by agricultural use in 1946. Production of DDT in the United States was greatest in 1959. Reduced prevalence of intersex in cricket frogs from 1960 to 1976 coincided with decreased use and the subsequent ban of most uses of DDT in the United States in 1972. Although exposure of larval tiger salamanders (*Ambystoma tigrinum*) to *p,p*-DDE (*p,p*-dichlorodiphenyl-dichloroethylene) stimulated growth of Mullerian ducts consistent with estrogenicity, exposure to technical-grade DDT had a paradoxical antiestrogenic effect (Clark et al. 1998). Thus, if cricket frogs responded to DDT exposure as did *A. tigrinum*, its use could have contributed to the concurrent decrease in female and increase in intersex cricket frogs during 1946–1959.

Atrazine was first marketed as a broadleaf herbicide for maize production in 1959, and use rapidly expanded. By 1993, the Midwest states of Illinois, Iowa, Nebraska, and Indiana accounted for 43% of the total amount of atrazine applied in the United States (Atrazine Ecological Risk Assessment Panel 1995), and it is still widely applied in the region. Based on findings of intersexuality at very low atrazine concentrations, Hayes et al. (2002, 2003) concluded that the widespread use of atrazine may have been a significant factor in amphibian declines. However, a recent study (Carr et al. 2003) indicated a higher threshold for atrazine-induced intersexuality in frogs. Additional research is needed to resolve this issue. Our study demonstrates that endocrine disruption and intersexuality were present in cricket frogs long before the advent of atrazine. However, the possibility that atrazine is one of the endocrine disruptors that contributed to the decline of cricket frogs and impedes expansion

of its populations in central and northern Illinois is not ruled out by our findings.

Although we observed a decrease in intersex cricket frogs from Illinois after 1946–1959, McCallum and Trauth (2003) noted a progressive increase in the proportion of cricket frogs in Arkansas with external developmental abnormalities during four time periods from 1957 to 2000. Johnson et al. (2003) reported an increased prevalence of parasite-induced malformations by the trematode *Ribeiroia* from 1946 to 2002. Different patterns of contaminant and trematode exposures, and responses of germinal gonadal versus somatic cells to teratogenic stimuli may be responsible for the contrasting trends. Collection efforts and methods were not recorded in the catalogs and would have varied over time. Records typically were limited to the collector's name and the location and date of acquisition in the field. Thus, bias that favored or limited collection of hermaphroditic cricket frogs cannot be completely ruled out. However, we have identified no basis for bias that would influence the likelihood of collecting intersex individuals. Hermaphroditic cricket frogs were collected by a wide range of investigators, and multiple hermaphrodites were found in the collections of many museums. Moreover, the museum records include no mention of behavioral or physical variation for any of the hermaphroditic specimens, and the collectors reported no knowledge of hermaphroditism. Finally, hermaphroditic individuals of this species can be identified only through gross dissection and histopathologic studies, which were not undertaken on these frogs prior to our research.

The geographic distribution of both endocrine disruption (intersexuality) and the decline of cricket frogs were congruent. The observed decline was evident after a period of sustained endocrine disruption, as indicated by a large increase in prevalence of intersex gonads and masculinization of the population. A plausible explanation for these observations is that exposures to antiestrogenic PAHs, PCBs, PCDFs, PCDDs, and DDT caused endocrine disruption, and this contributed to the virtual disappearance of cricket frogs from the Chicago region. The intersex prevalence in cricket frogs in recent years is low and may represent a near-baseline condition. A suite of endocrine-disrupting organochlorine contaminants persists in soils and waters of the Midwest, but at substantially reduced levels compared with earlier decades (Abramowicz 1990). However, we cannot conclude that the era of endocrine disruption in cricket frogs has come to an end, because in areas with the most severe decline in populations and most severe endocrine disruption historically, numbers of remaining cricket frogs are now insufficient to permit sampling.

REFERENCES

- Abramowicz DA. 1990. Aerobic and anaerobic biodegradation of PCBs: a review. *Biotechnology* 10:241–251.
- Alford RA, Dixon PM, Pechmann JHK. 2001. Ecology: Global amphibian population declines. *Nature* 412:499–500.
- Atrazine Ecological Risk Assessment Panel. 1995. Ecological Risk Assessment of Atrazine in North American Surface Water. Greensboro, NC:Ciba Crop Protection.
- ATSDR. 1995. Toxicological Profile for Polycyclic Aromatic Hydrocarbons. Atlanta, GA:Agency for Toxic Substances and Disease Registry. Available: <http://www.atsdr.cdc.gov/toxprofiles/tp69.html> [accessed 14 January 2005].
- Baatrup E, Junge M. 2001. Antiandrogenic pesticides disrupt sexual characteristics in the adult male guppy (*Poecilia reticulata*). *Environ Health Perspect* 109:1063–1070.
- Bergeron JM, Crews D, McLachlan JA. 1994. PCBs as environmental estrogens: turtle sex determination as a biomarker of environmental contamination. *Environ Health Perspect* 102:780–781.
- Blaustein AR, Wake DB. 1990. Declining amphibian populations: a global phenomenon? *Trends Ecol Evol* 5:203–204.
- Brodman R, Kilmer M. 1998. Status of amphibians in northwestern Indiana. In: *Status and Conservation of Midwestern Amphibians* (Lannoo MJ, ed). Iowa City, IA:University of Iowa Press, 125–136.
- Carr JA, Gentles A, Smith EE, Goleman WL, Urquidí LJ, Thuet K, et al. 2003. Response of larval *Xenopus laevis* to atrazine: assessment of growth, metamorphosis, and gonadal and laryngeal morphology. *Environ Toxicol Chem* 22:396–405.
- Caspar GS. 1998. Review of the status of Wisconsin amphibians. In: *Status and Conservation of Midwestern Amphibians* (Lannoo MJ, ed). Iowa City, IA:University of Iowa Press, 199–205.
- Chaloupka K, Harper N, Krishnan V, Santostefano M, Rodriguez LV, Safe S. 1993. Synergistic activity of polynuclear aromatic hydrocarbon mixtures as aryl hydrocarbon (Ah) receptor agonists. *Chem Biol Interact* 89:141–158.
- Chaloupka K, Krishnan V, Safe S. 1992. Polynuclear aromatic hydrocarbon carcinogens as antiestrogens in MCF-7 human breast cancer cells: role of the Ah receptor. *Carcinogenesis* 13:2233–2239.
- Clark EJ, Norris DO, Jones RE. 1998. Interactions of gonadal steroids and pesticides (DDT, DDE) on gonaduct growth of larval tiger salamanders, *Ambystoma tigrinum*. *Gen Comp Endocrinol* 109:94–105.
- Diana SG, Resetaritis WJ Jr, Schaeffer DJ, Beckmen KB, Beasley VR. 2000. Effects of atrazine on amphibian growth and survival in artificial aquatic communities. *Environ Toxicol Chem* 19:2961–2967.
- Fry DM, Toone CK. 1981. DDT-induced feminization of gull embryos. *Science* 213:922–924.
- Guillette LJ Jr, Crain DA, Rooney AA, Pickford DB. 1995. Organization versus activation: the role of endocrine disrupting contaminants (EDCs) during embryonic development in wildlife. *Environ Health Perspect* 103(suppl 7):157–164.
- Guillette LJ Jr, Gross TS, Masson GR, Matter JM, Percival HF, Woodward AR. 1994. Developmental abnormalities of the gonad and abnormal sex hormone concentrations in juvenile alligators from contaminated and control lakes in Florida. *Environ Health Perspect* 102:680–688.
- Hansen L. 1987. Environmental toxicology of polychlorinated biphenyls. In: *Environmental Toxin Series 1: Polychlorinated Biphenyls (PCBs): Mammalian and Environmental Toxicology* (Safe S, Hutzinger O, eds). Vol 1. New York:Springer-Verlag, 15–48.
- Hayes T, Haston K, Tsui M, Hoang A, Haeffle C, Vonk A. 2003. Atrazine-induced hermaphroditism at 0.1 ppb in American leopard frogs (*Rana pipiens*): laboratory and field evidence. *Environ Health Perspect* 111:568–575.
- Hayes TB, Collins A, Lee M, Mendoza M, Noriega N, Stuart AA, et al. 2002. Hermaphroditic, demasculinized frogs after exposure to the herbicide atrazine at low ecologically relevant doses. *Proc Natl Acad Sci USA* 99:5476–5480.
- Houlahan JE, Findlay CS, Schmidt BR, Meyer AH, Kuzman SL. 2000. Quantitative evidence for global amphibian population declines. *Nature* 404:752–755.
- Jansen HT, Cook PS, Porcelli J, Liu TC, Hansen LG. 1993. Estrogenic and antiestrogenic actions of PCBs in the female rat: *in vitro* and *in vivo* studies. *Reprod Toxicol* 7:237–248.
- Johnson PT, Lunde KB, Zelmer DA, Werner JK. 2003. Limb

- deformities as an emerging parasitic disease in amphibians: evidence from museum specimens and resurvey data. *Conserv Biol* 17:1724–1737.
- Jung RE. 1993. Blanchard's cricket frogs (*Acris crepitans blanchardi*) in southwest Wisconsin. *Tran Wisc Acad Sci Arts Lett* 81:79–87.
- Krishnan V, Safe S. 1993. Polychlorinated biphenyls (PCBs), dibenzo-*p*-dioxins (PCDDs), and dibenzofurans (PCDFs) as antiestrogens in MCF-7 human breast cancer cells: quantitative structure-activity relationships. *Toxicol Appl Pharmacol* 120:55–61.
- Li MH, Hansen LG. 1996. Enzyme induction and acute endocrine effects in prepubertal female rats receiving environmental PCB/PCDF/PCDD mixtures. *Environ Health Perspect* 104:712–722.
- Li MH, Zhao YD, Hansen LG. 1994. Multiple dose toxicokinetic influence on the estrogenicity of 2,2',4,4',5,5'-hexachlorobiphenyl. *Bull Environ Contam Toxicol* 53:583–590.
- Ludwig DR, Redmer M, Domazlicky R, Kobal S, Conklin B. 1992. Current status of amphibians and reptiles in DuPage County, Illinois. *Trans Ill St Acad Sci* 85:187–199.
- Machala M, Ciganek M, Blaha L, Minksova K, Vondrak J. 2001. Aryl hydrocarbon receptor-mediated and estrogenic activities of oxygenated polycyclic aromatic hydrocarbons and azaarenes originally identified in extracts of river sediments. *Environ Toxicol Chem* 20:2736–2743.
- Malby TA, Bjerke DL, Moore RW, Fendron-Fitzpatrick A, Peterson RE. 1992. *In utero* and lactational exposure of male rats to 2,3,7,8-tetrachlorodibenzo-*p*-dioxin. 3. Effects on spermatogenesis and reproductive capability. *Toxicol Appl Pharmacol* 114:118–126.
- Matta MB, Cairncross C, Kocan RM. 1998. Possible effects of polychlorinated biphenyls on sex determination in rainbow trout. *Environ Toxicol Chem* 17:26–39.
- McCallum ML, Trauth SE. 2003. A forty-three year museum study of northern cricket frog (*Acris crepitans*) abnormalities in Arkansas: upward trends and distributions. *J Wildl Dis* 39:522–528.
- Mellanby K. 1992. *The DDT Story*. Farnham, England: British Crop Protection Council.
- Metcalfe TL, Metcalfe CD, Kiparissis Y, Niimi AJ, Foran CM, Benson WH. 2000. Gonadal development and endocrine responses in Japanese medaka (*Oryzias latipes*) exposed to *o,p'*-DDT in water or through maternal material. *Environ Toxicol Chem* 19:1893–1900.
- Mierzwa KS. 1998. Status of northeastern Illinois amphibians. In: *Status and Conservation of Midwestern Amphibians* (Lannoo MJ, ed). Iowa City, IA: University of Iowa Press, 115–124.
- Minton SA. 1998. Observations on Indiana amphibian populations: a forty-five-year overview. In: *Status and Conservation of Midwestern Amphibians* (Lannoo MJ, ed). Iowa City, IA: University of Iowa Press, 217–220.
- Moriarty JJ. 1998. Status of amphibians in Minnesota. In: *Status and Conservation of Midwestern Amphibians* (Lannoo MJ, ed). Iowa City, IA: University of Iowa Press, 166–168.
- Mossman MJ, Hartman LM, Hay R, Sauer JR, Dhuey BJ. 1998. Monitoring long-term trends in Wisconsin frog and toad populations. In: *Status and Conservation of Midwestern Amphibians* (Lannoo MJ, ed). Iowa City, IA: University of Iowa Press, 169–198.
- Palmer BD, Huth LK, Pioto DL, Selcer KW. 1998. Vitellogenin as a biomarker for xenobiotic estrogens in an amphibian model system. *Environ Toxicol Chem* 17:30–36.
- Palmer BD, Palmer SK. 1995. Vitellogenin induction by xenobiotic estrogens in the red-eared slider turtle and African clawed frog. *Environ Health Perspect* 103(suppl 4):19–25.
- Qin ZF, Zhou JM, Chu SG, Xu XB. 2003. Effects of Chinese domestic polychlorinated biphenyls (PCBs) on gonadal differentiation in *Xenopus laevis*. *Environ Health Perspect* 111:553–556.
- Reeder AL, Foley G, Nichols D, Wikoff B, Faeh S, Eisold J, et al. 1998. Forms and prevalence of intersexuality and effects of environmental contaminants on sexuality in cricket frogs (*Acris crepitans*). *Environ Health Perspect* 106:261–266.
- Shaffer HB, Fisher RN, Davidson C. 1998. The role of natural history collections in documenting species declines. *Trends Ecol Evol* 13:27–30.
- Smith PW. 1961. *The Amphibians and Reptiles of Illinois*. Urbana, IL: Illinois Natural History Survey.
- Sullivan KB, Spence KM. 2003. Effects of sublethal concentrations of atrazine and nitrate on metamorphosis of the African clawed frog. *Environ Toxicol Chem* 22:627–635.
- Thomas P, Budiantara L. 1995. Reproductive life history stages sensitive to oil and naphthalene in Atlantic croaker. *Marine Environ Res* 39:147–150.
- Tyler MJ. 1991. Declining amphibian populations—a global phenomenon? An Australian perspective. *Alytes* 9:43–50.
- U.S. Army Service Forces. 1946. *Application of DDT by Airplane for Mosquito Control at Savanna Ordnance Depot Proving Grounds, Illinois, 1946*. Chicago: Army Service Forces, Sixth Service Command, Repairs and Utilities Division.
- Vogt RC. 1981. *Natural History of Amphibians and Reptiles in Wisconsin*. Milwaukee, WI: Milwaukee Public Museum.